

## FREE PRODUCTS OF LIE GROUPS

BY

SIDNEY A. MORRIS (SYDNEY)

**1. Introduction.** In [1] Graev proved that the algebraic free product of two Hausdorff topological groups can be equipped with a topology making it into a Hausdorff topological group and satisfying the appropriate conditions for a coproduct in the category of topological groups; see also [2] and [5].

In [3] and [4] we showed that a free product of connected locally compact groups is never locally compact. We prove here that a free product of connected locally compact groups  $G_1$  and  $G_2$  has no small subgroups if and only if  $G_1$  and  $G_2$  have no small subgroups (that is,  $G_1$  and  $G_2$  are Lie groups). A similar result is true if  $G_1$  or  $G_2$  is only assumed to be compact.

**2. Preliminaries.**

**Definition 1.** Let  $G_1$  and  $G_2$  be topological groups. Then the topological group  $F$  is said to be a *free product* of  $G_1$  and  $G_2$ , denoted by  $G_1 * G_2$ , if it has the following properties:

(a)  $G_1$  and  $G_2$  are subgroups of  $F$ .

(b)  $F$  is generated algebraically by  $G_1 \cup G_2$ .

(c) If  $\gamma_1$  and  $\gamma_2$  are continuous homomorphisms of  $G_1$  and  $G_2$ , respectively, into a topological group  $H$ , then there exists a continuous homomorphism  $\Gamma$  of  $F$  into  $H$  such that  $\Gamma = \gamma_i$  on  $G_i$  for  $i = 1, 2$ .

Of course a free product is simply the coproduct in the category of all topological groups. Thus, if it exists, it is unique up to isomorphism. Free products of topological groups were first studied by Graev [1]. He showed that the free product of Hausdorff groups exists and is Hausdorff. His proof showed that the underlying group structure of  $G_1 * G_2$  is the algebraic free product of  $G_1$  and  $G_2$ .

**Definition 2.** Let  $G$  be a group and  $X$  a subset which generates  $G$  algebraically. Then  $a \in G$  is said to be of *length  $n$  with respect to  $X$*  if  $n$  is the least integer  $N$  such that  $a = x_1^{\varepsilon_1} \dots x_N^{\varepsilon_N}$ , where  $\varepsilon_i = \pm 1$  and  $x_i \in X$ .

for  $i = 1, \dots, N$ . The set of all elements in  $G$  of length not greater than  $m$  will be denoted by  $G_m(X)$ .

Our first preliminary theorem is essentially proved in [4].

**THEOREM A.** For  $i = 1, 2$ , let  $G^i$  be a Hausdorff group with a compact subspace  $X_i$  which generates  $G^i$  algebraically. Further, let the topology of  $G^i$  be the finest group topology on  $G^i$  which induces the same topology on  $X_i$ . If  $G = G^1 * G^2$  is the free product of  $G^1$  and  $G^2$ , then  $X = X_1 \cup X_2$  is a compact set which generates  $G$  algebraically and has the property that a subset  $V$  of  $G$  is closed if and only if  $V \cap G_n(X)$  is compact, for each  $n$ .

**THEOREM B** (see [4]). If  $G$  is a connected locally compact group, then there exists a compact subset  $X$  of  $G$  such that

- (i)  $X$  generates  $G$  algebraically,
- (ii) the topology of  $G$  is the finest group topology on  $G$  which induces the given topology on  $X$ .

**Notation.** We will use  $e$  to denote the identity element of a group.

Whenever we write  $x_1 x_2 \dots x_n \in G_1 * G_2$  we will mean that, for any  $i$ ,  $x_i \neq e$  and  $x_i$  and  $x_{i+1}$  are not both in  $G_1$  or  $G_2$ .

## 2. Results.

**THEOREM 1.** Let  $G^1$  and  $G^2$  be connected locally compact groups. Then  $G = G^1 * G^2$  has no small subgroups if and only if  $G^1$  and  $G^2$  have no small subgroups.

**Proof.** If  $G^1 * G^2$  has no small subgroups, then  $G^1$  and  $G^2$ , being subgroups of  $G^1 * G^2$ , have no small subgroups.

Conversely, assume that  $G^1$  and  $G^2$  have no small subgroups; that is, they are Lie groups. Let  $O_1$  and  $O_2$  be open neighbourhoods of  $e$  in  $G^1$  and  $G^2$ , respectively, which contain no non-trivial subgroups.

For  $i = 1, 2$ , by Theorem B, we can find a compact subset  $X_i$  of  $G^i$  which generates  $G^i$  algebraically and is such that the topology of  $G^i$  is the finest group topology on  $G^i$  which induces the same topology on  $X_i$ . Put  $X = X_1 \cup X_2$ . Then, by Theorem A,  $X$  generates  $G$  algebraically and a subset  $C$  of  $G$  is closed if and only if  $C \cap G_n(X)$  is compact for every  $n$ .

Let  $d$  be a compatible metric on  $G^1$  and define

$$A_n = G_n^1(X_1) \cap \left\{ x: x \in G^1 \text{ and } d(x, e) \geq \frac{1}{n} \right\} \quad \text{for each } n \geq 1.$$

Then  $A_n$  is compact.

Similarly define compact subsets  $B_n$  of  $G^2$ . We note that

$$(1) \quad A_n \supseteq A_{n-1} \text{ and } B_n \supseteq B_{n-1} \quad \text{for every } n \geq 2$$

and

$$(2) \quad \bigcup_{n=1}^{\infty} A_n = G^1 - \{e\} \quad \text{and} \quad \bigcup_{n=1}^{\infty} B_n = G^2 - \{e\}.$$

Define

$$V_1 = A_1 \cup B_1, \quad V_2 = A_2 B_2 \cup B_2 A_2, \quad V_3 = A_3 B_3 A_3 \cup B_3 A_3 B_3$$

and so on. Clearly, each  $V_n$  is compact.

For  $n = 0, 1, 2, \dots$  and  $m = 1, 2, \dots$ , define

$$W_n^m = \{(x_1 \dots x_n)^{-1} y x_1 \dots x_n : \text{each } x_i \in G_m(X) \text{ and } y \in [A_m \cap (G^1 - O_1)] \cup [B_m \cap (G^2 - O_2)]\}.$$

We note that each  $W_n^m$  is compact; every element in  $W_n^m$  has length, with respect to  $X$ , greater than  $n$ .

Without loss of generality assume that there is an integer  $N$  such that

$$O_i \subseteq \left\{ x : x \in G^i \text{ and } d(x, e) \leq \frac{1}{N} \right\}, \quad i = 1, 2.$$

Then, for  $m > k > N$ ,

$$A_m \cap (G^1 - O_1) \cap G_k(X) = A_k \cap (G^1 - O_1) \cap G_k(X).$$

Thus, for  $m > k > N$  and any  $n > 0$ ,

$$W_n^m \cap G_k(X) = W_n^k \cap G_k(X).$$

Our comments on the sets  $W_n^m$  show that if

$$V_0 = \bigcup_{n=0}^{\infty} \bigcup_{m=1}^{\infty} W_n^m,$$

then

$$V_0 \cap G_k(X) = \left[ \bigcup_{n=0}^k \bigcup_{m=1}^k W_n^m \right] \cap G_k(X) \quad \text{for any } k > N.$$

Thus  $V_0 \cap G_k(X)$  is compact for every  $k \geq 1$ . By Theorem A, this implies that  $V_0$  is closed in  $G$ .

We now define

$$V = \bigcup_{n=0}^{\infty} V_n.$$

For any  $k$  we have

$$V \cap G_k(X) = [V_0 \cap G_k(X)] \cup \left[ \bigcup_{n=1}^{\infty} V_n \cap G_k(X) \right].$$

Since each element in  $V_n$ ,  $n \geq 1$ , has the length greater than or equal to  $n$ , we see that

$$V \cap G_k(X) = [V_0 \cap G_k(X)] \cup \left[ \bigcup_{n=1}^k V_n \cap G_k(X) \right].$$

Since  $V_0$  is closed in  $G$  and each  $V_n$ ,  $n \geq 1$ , is compact, we infer that  $V \cap G_k(X)$  is compact for all  $k$ . Thus, by Theorem A,  $V$  is closed in  $G$ . Further,  $e \notin V$ . Thus  $G - V$  is an open neighbourhood of  $e$ .

To complete the proof we only have to show that

(\*) For any  $a \in G$  with  $a \neq e$  we have  $a^k \in V$  for some  $k$ .

Suppose  $a \in G$  and  $a \neq e$ . Then  $a = x_1 \dots x_n$ . It follows from (1) and (2) that there exists an  $m$  such that  $x_i \in A_m \cup B_m$  for  $i = 1, \dots, n$ .

Case 1. The length of  $a^2$  is greater than the length of  $a$ . Then there exists a  $k$  such that  $a^k$  has the length greater than  $m$ . Clearly,  $a^k \in V_r$ , where  $r$  denotes the length of  $a^k$ , since  $a^k$  is in the group generated by  $\{x_1, \dots, x_n\}$  and each  $x_i \in A_r \cup B_r$ . So  $a^k \in V$ , as required.

Case 2. The length of  $a^2$  is less than or equal to the length of  $a$ . This can happen only if  $a = (z_1 \dots z_t)^{-1} y z_1 \dots z_t$  for some  $y \in G^1 \cup G^2$  and  $t \geq 0$ . Without loss of generality assume  $y \in G^1$ . Since  $O_1$  contains no non-trivial subgroup, there exists a  $k \geq 1$  such that  $y^k \notin O_1$ . By (2), there exists an  $m$  such that  $y^k \in A_m$ . Thus

$$a^k = (z_1 \dots z_t)^{-1} y^k z_1 \dots z_t \in V_0.$$

Hence  $a^k \in V$  and the proof of (\*) is complete.

In a similar manner we can prove

**THEOREM 2.** *Let  $G^1$  and  $G^2$  be locally compact groups which are either connected or compact. Then  $G = G^1 * G^2$  has no small subgroups if and only if  $G^1$  and  $G^2$  have no small subgroups.*

#### REFERENCES

- [1] М. И. Граев, *О свободных произведениях топологических групп*, Известия Академии Наук СССР, Серия математическая, 14 (1950), p. 343-354.
- [2] S. A. Morris, *Free products of topological groups*, Bulletin of the Australian Mathematical Society 4 (1971), p. 17-29.
- [3] — *Local compactness and free products of topological groups*, Proceedings of the Royal Society of New South Wales (Australia), to appear.
- [4] — *Local compactness and local invariance of free products of topological groups* (submitted for publication).
- [5] E. T. Ordman, *Free products of topological groups with equal uniformities, I*, Colloquium Mathematicum 31 (1974), p. 37-43.

UNIVERSITY OF NEW SOUTH WALES

*Reçu par la Rédaction le 10. 8. 1973*