

FREE PRODUCTS OF TOPOLOGICAL GROUPS WITH AMALGAMATION

ELYAHU KATZ AND SIDNEY A. MORRIS

It is proved that the free product of any two k_ω -groups with a compact subgroup amalgamated is a k_ω -group, and in particular, Hausdorff.

1. Introduction. In recent years much work has been done on describing the topology of free products of topological groups (see for example [1, 3, 8, 10, 12, 13, 15]). From there it is natural to progress to free products with amalgamation.

One would hope that the free product with amalgamation of any Hausdorff topological groups exists, is Hausdorff and its underlying group structure is the amalgamated free product of the underlying groups. This would include as a special case Graev's theorem [2] that the free product of Hausdorff groups is Hausdorff. As his proof is certainly non-trivial, it should not be expected that this "hope" will be easily verified, even if the result is true.

The first contribution to this problem was by Ordman [13], who showed that the amalgamated free product of certain locally invariant Hausdorff topological groups is Hausdorff. The next contribution was by Khan and Morris [5] who proved the Hausdorffness of the free product of Hausdorff groups with a central subgroup amalgamated. This has recently been extended by Katz and Morris [4] to free products of k_ω -groups with a closed normal subgroup amalgamated.

Most of the work on free topological groups and free products of topological groups in fact deals with topological groups which are k_ω -spaces. Therefore, the result we would like to have is that the amalgamated free product of k_ω -groups is a k_ω -group. This would imply La Martin's theorem that epics in the category of k_ω -groups have dense range. (See [6], [11] and [14].) We prove here that the free product of any two k_ω -groups with a compact subgroup amalgamated is a k_ω -group and in particular Hausdorff. This result includes a large class of examples not covered by [4] or [13], since every connected locally compact Hausdorff topological group, G is a k_ω -group and has a compact subgroup K such

that G is homeomorphic to $\mathbf{R}^n \times K$, for some Euclidean group \mathbf{R}^n . Such examples with G not locally invariant and K not normal exist in profusion.

2. Definitions and statements of the main result. The standard references for amalgamated free products of groups are B. H. Neumann [9] and Magnus, Karrass and Solitar [7]. For completeness we include some definitions here.

DEFINITION. Let A be a common subgroup of groups G and H . The group $G *_A H$ is said to be the *free product of G and H with amalgamated subgroup A* if

- (i) G and H are subgroups of $G *_A H$,
- (ii) $G \cup H$ generates $G *_A H$ algebraically,
- (iii) every pair ϕ_1, ϕ_2 of homomorphisms of G and H , respectively, into any group D which agree on A , extend to a homomorphism Φ of $G *_A H$ into D .

DEFINITION. Let A be a common subgroup of topological groups G and H . The topological group $G *_A H$ is said to be the *free product of the topological groups G and H with amalgamated subgroup A* if

- (i) G and H are topological subgroups of $G *_A H$,
- (ii) $G \cup H$ generates $G *_A H$ algebraically, and
- (iii) every pair ϕ_1, ϕ_2 of continuous homomorphisms of G and H , respectively, into any topological group D , which agree on A , extend to a continuous homomorphism of $G *_A H$ into D .

Our main result is the following:

THEOREM. *Let F and G be k_ω -groups with a common compact subgroup A . Then the free product of F and G with A amalgamated is a k_ω -group. (In particular, the amalgamated free product $F *_A G$ is Hausdorff.)*

3. Yoffi representations. To prove the main theorem it suffices to show that the kernel, K , of the canonical homomorphism $\Phi: F * G \rightarrow F *_A G$ is closed. The first step in doing this is to find a nice representation for the elements of K . This will be achieved in Proposition 1.

NOTATION. We denote the embedding map of A in F by f and the embedding map of A in G by g .

It is readily seen that K is the normal subgroup generated by $\{f(a)g(a)^{-1}: a \in A\}$. Further, observe that each $w \in K$ has a representation.

$$(*) \quad \begin{cases} w = (t_{1,1} \cdots t_{1,q_1} v_1 t_{1,q_1}^{-1} \cdots t_{1,1}^{-1}) (t_{2,1} \cdots t_{2,q_2} v_2 t_{2,q_2}^{-1} \cdots t_{2,1}^{-1}) \\ \cdots (t_{s,1} \cdots t_{s,q_s} v_s t_{s,q_s}^{-1} \cdots t_{s,1}^{-1}), \end{cases}$$

where for each j , $t_{j,i}$ lies in F or G , and $t_{j,i}$ and $t_{j,i+1}$ do not both belong to F or both belong to G , and each $v_i \in \{f(a)g(a)^{-1}, g(a)f(a)^{-1}: a \in A\}$.

DEFINITIONS. For each representation of the form $(*)$ we define the *weight of the representation* to be a sequence $(a_1, a_2, a_3, \dots, a_n, \dots)$, where a_i is the number of q 's equal to the number i . We well-order the set of sequences of saying $(a_1, a_2, \dots, a_n, \dots) < (a'_1, a'_2, \dots, a'_n, \dots)$ if and only if $a_i = a'_i$, for $i > n$ and $a_n < a'_n$.

Given any word w it may have many representations of the form $(*)$. We define the *weight of the word* w to be the least weight of all the representations of w .

We shall see later, when dealing with transpositions, that a word need not have a unique representation of least weight.

REMARK 1. Let w have representation $(*)$ of least weight.

(i) If $v_i = f(a_i)g(a_i)^{-1}$, then $t_{i,q_i} \neq f(a_i)^{-1}$ and $t_{i,q_i}^{-1} \neq g(a_i)$, since otherwise w would have a representation of smaller weight.

(ii) No $t_{i,1} \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}$ equals $x_1 x_2 \cdots x_r$ where each $x_j = s_{j,1} \cdots s_{j,n_j} f(b_j) g(b_j)^{-1} s_{j,n_j}^{-1} \cdots s_{j,1}^{-1}$, where each s_i lies in F or G and s_i and s_{i+1} do not both belong to F or both belong to G and $b_j \in A$, for $n_j < q_i, j = 1, 2, \dots, r$. (This would contradict the fact that the representation of w has least weight.)

(iii) If

$$\begin{aligned} & (t_{i,1} \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}) (t_{(i+1),1} \cdots t_{(i+1),q_{i+1}} v_{i+1} t_{(i+1),q_{i+1}}^{-1} \cdots t_{(i+1),1}^{-1}) \\ & \cdots (t_{(i+h),1} \cdots t_{(i+h),q_{i+h}} v_{i+h} t_{(i+h),q_{i+h}}^{-1} \cdots t_{(i+h),1}^{-1}) \\ & = s_1 \cdots s_q v_s q^{-1} \cdots s_1^{-1}, \end{aligned}$$

where $v = f(a)g(a)^{-1}$, $a \in A$, and each $s_i \in F$ or G then $q > \max\{q_i, q_{i+1}, \dots, q_{i+h}\}$.

(iv) Let

$$x_i = t_{i,1} \cdots t_{i,q_i} g(a_i) f(a_i)^{-1} t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}.$$

Put $a = a_i^{-1}$. If $t_{i,q_i} \in F$, put $t = t_{i,q_i} f(a_i)$, so $t \in F$. If $t_{i,q_i} \in G$, put $t = t_{i,q_i} g(a_i)$, so $t \in G$. Then in both cases

$$x_i = t_{i,1} \cdots t_{i,q_{i-1}} t f(a) g(a)^{-1} t_{i,q_{i-1}}^{-1} \cdots t_{i,1}^{-1}.$$

The above rewriting shows that we can assume that if $(*)$ is a representation of a word $w \in K$, then each v_i is of the form $f(a_i)g(a_i)^{-1}$. Because, if it is not, we can change the representation so that it is. Further, the change of representation does not affect the weight. So each $w \in K$ has a representation of the form $(*)$ of least weight, with each $v_i = f(a_i)g(a_i)^{-1}$, for some $a_i \in A$.

NOTATION. From now on whenever we say we have a representation of the form $(*)$ we mean that $v_i = f(a_i)g(a_i)^{-1}$, for each i .

We need some further restrictions on the representations we consider.

DEFINITIONS. Let $x_1 = t_{1,1} \cdots t_{1,q_1} f(a_1)g(a_1)^{-1} t_{1,q_1}^{-1} \cdots t_{1,1}^{-1}$ and $x_2 = t_{2,1} \cdots t_{2,q_2} f(a_2)g(a_2)^{-1} t_{2,q_2}^{-1} \cdots t_{2,1}^{-1}$.

(i) Let $q_1 \geq q_2$ and $t_{1,j} = t_{2,j}$ for $1 \leq j \leq q_2 - 1$ and $t_{1,q_2}^{-1} t_{2,q_2} f(a_2) = e$. If $q_1 = q_2$ then x_1 is said to have an *even link* with x_2 . If $q_1 \neq q_2$ then x_1 is said to have an *odd link* with x_2 .

(ii) Let $q_1 \leq q_2$ and $t_{1,j} = t_{2,j}$ for $1 \leq j \leq q - 1$ and $g(a_1)^{-1} t_{1,q_1}^{-1} t_{2,q_1} = e$. If $q_1 = q_2$ then x_1 is said to have an *even link* with x_2 . If $q_1 \neq q_2$ then x_1 is said to have an *odd link* with x_2 .

(iii) If x_1 does not have an even link with x_2 or an odd link with x_2 , then x_1 is said to have a *neutral link* with x_2 .

(iv) A representation $x_1 x_2 \cdots x_s$ for $w \in K$, where each $x_i = t_{i,1} \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}$, for $v_i = f(a_i)g(a_i)^{-1}$, is said to be *yoffi*¹ if it has least weight and each x_i has a neutral link with x_{i+1} .

LEMMA 1. *Let the representation $(*)$ of $w \in K$ have least weight and be such that $q_i = q_{i+1}$ for some i . If $t_{i,j} = t_{i+1,j}$, $j = 1, \dots, (q_i - 1)$, $v_i = f(a_i)g(a_i)^{-1}$ and $v_{i+1} = f(a_{i+1})g(a_{i+1})^{-1}$ then $t_{i,q_i}^{-1} t_{i+1,q_{i+1}} f(a_{i+1}) \neq e$ and $g(a_i)^{-1} t_{i,q_i}^{-1} t_{i+1,q_{i+1}} \neq e$; that is, x_i does not have an even link with x_{i+1} .*

Proof. Suppose $g(a_i)^{-1} t_{i,q_i}^{-1} t_{i+1,q_{i+1}} = e$. Then

$$(t_{i+1,q_{i+1}})^{-1} = g(a_i)^{-1} t_{i,q_i}^{-1}.$$

So $[t_{i,1} \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}]$.

¹A Hebrew word meaning *nice*.

Proof. Choose a representation of w of the form $(*)$ which has least weight. Let this representation be $w = x_1 \cdots x_s$, where $x_i = t_{i,1} \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}$ where $v_i = f(a_i)g(a_i)^{-1}$ for $i = 1, \dots, s$.

Our proof will be by induction on s . Firstly consider the case $s = 2$; that is, $w = x_1 \cdot x_2$. If x_1 has a neutral link with x_2 , then this is a yoffi representation. Observe that by Lemma 1, x_1 cannot have an even link with x_2 . So we let x_1 have an odd link with x_2 . Then by Lemma 2 we can apply a transposition to obtain $x'_2 x'_1$ such that this representation also has least weight and x'_2 has a neutral link with x'_1 . Hence $x'_2 x'_1$ is a yoffi representation of w .

Now suppose that every word of the form $w = x_1 x_2 \cdots x_n$, for $n \leq s$, has a yoffi representation. Then consider any word

$$w = x_1 x_2 \cdots x_{s+1}$$

where the weight of the word w equals the weight of the representation $x_1 \cdots x_{s+1}$. Clearly the weight of the word $x_1 x_2 \cdots x_s$ equals the weight of the representations $x_1 \cdots x_s$. Thus by the inductive hypothesis it has a yoffi representation of the same weight. So without loss of generality we can assume that it is $x_1 \cdots x_s$. So each x_i has a neutral link with x_{i+1} for $i = 1, \dots, s-1$. If x_s has a neutral link with x_{s+1} we are done. If not, we perform a transposition of $x_s x_{s+1}$ to obtain from w the representation $x_1 x_2 \cdots x_{s-1} x'_{s+1} x'_s$. If x_{s-1} has a neutral link with x'_{s+1} , we are done. If x_{s-1} has an odd link with x'_{s+1} then we apply a transposition of $x_{s-1} x'_{s+1}$ to obtain the representation $x_1 x_2 \cdots x_{s-2} x''_{s+1} x'_{s-1} x'_s$. By Lemma 3 (proved independently below) $x'_{s-1} x'_s$ has a neutral link. Once again, if $x_{s-2} x''_{s+1}$ has a neutral link we are done. If not, apply a transposition of $x_{s-2} x''_{s+1}$ to obtain $x_1 x_2 \cdots x_{s-3} x'''_{s+1} x'_{s-2} x'_{s-1} x'_s$. By Lemma 3 again $x'_{s-2} x'_s$ has a neutral link. Continuing this process, if necessary, we obtain a yoffi representation. So the proof is complete once we have Lemma 3.

LEMMA 3. *Let $x_1 x_2 x_3$ be a representation of a word w such that the weight of the word w equals the weight of the representation $x_1 x_2 x_3$. If x_1 has a neutral link with x_2 , then $x_1 x_2 x_3$, $x_1 x'_3 x'_2$ or $x'_3 x'_1 x'_2$ is a yoffi representation where $x'_3 x'_2$ is a transposition of $x_2 x_3$ and $x'_3 x'_1$ is a transposition of $x_1 x'_3$.*

Proof. If $x_1 x_2 x_3$ is not a yoffi representation let

$$x_i = t_{i,1} \cdots t_{i,q_i} f(a_i) g(a_i)^{-1} t_{i,1}^{-1} \cdots t_{i,1}^{-1}, \quad \text{for } i = 1, 2, 3.$$

We distinguish six cases:

- (i) $q_1 < q_2 < q_3$, (ii) $q_1 < q_3 < q_2$, (iii) $q_3 < q_2 < q_1$,
- (iv) $q_2 < q_3 < q_1$, (v) $q_3 < q_1 < q_2$, (vi) $q_2 < q_1 < q_3$.

Consider case (i). We claim that x_1 has a neutral link with x'_3 since x_1 has a neutral link with x_2 and the first q_1 elements of x_2 and x'_3 are the same. Hence $x_1x'_3x'_2$ is a yoffi representation. Exactly the same argument works in case (ii). Now consider case (iii). If x_1 has a neutral link with x'_3 we are done. If not, consider $x''_3x'_1x'_2$. We have to show that x'_1 has a neutral link with x'_2 . In obtaining x'_2 from x_2 we replace only t_{2,q_3} by t_{3,q_3} and t_{2,q_3+1} by $g(a_3)t_{2,q_3+1}$ and in obtaining x'_1 from x_1 we replace only t_{1,q_3} by t_{3,q_3} and t_{1,q_3+1} by $g(a_3)t_{1,q_3+1}$, because $x_3 = x'_3$ in this case. The fact that x_1 has a neutral link with x_2 then clearly implies that x'_1 has a neutral link with x'_2 . Hence $x''_3x'_1x'_2$ is a yoffi representation.

Cases (iv)–(vi) are proved in the same fashion.

Finally observe that $q_2 = q_3$ implies that x_2 has a neutral link with x_3 and so $x_1x_2x_3$ is a yoffi representation, $q_1 = q_2$ implies $x''_3x'_1x'_2$ is a yoffi representation, and $q_1 = q_3$ implies $x_1x'_3x'_2$ is a yoffi representation. This completes the proof of the lemma.

4. The reduced lengths of elements of K . The key results of this section are Propositions 2 and 3 which imply that if $w = x_1x_2 \cdots x_s$ is a yoffi representation and w has reduced length $\leq n$ (that is, $w \in (F \cup G)^n$), then $s \leq n$ and $2q_i + 1 \leq 4n$, for $i = 1, 2, \dots, s$.

LEMMA 4. *Let $w \in K$ have representation (*) of least weight. If $q_i > q_{i+1}$ for some i and $t_{i,j} = t_{i+1,j}$, for $j = 1, \dots, q_{i+1}$, then $t_{i(q_{i+1}+1)} \neq f(a_{i+1})$.*

Proof. Suppose $t_{i(q_{i+1}+1)} = f(a_{i+1})$. Then

$$\begin{aligned} & t_{i,1} \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \cdots t_{i,1}^{-1} \\ &= \left(t_{i,1} \cdots t_{i,q_{i+1}} f(a_{i+1}) g(a_{i+1})^{-1} t_{i,q_{i+1}}^{-1} \cdots t_{i,1}^{-1} \right) \\ & \quad \cdot \left(t_{i,1} \cdots \left[t_{i,q_{i+1}} g(a_{i+1}) t_{i(q_{i+1}+2)} \right] \cdots t_{i,q_i} v_i t_{i,q_i}^{-1} \right. \\ & \quad \quad \quad \left. \cdots \left[t_{i,q_{i+1}+2}^{-1} g(a_{i+1})^{-1} t_{i,q_{i+1}+1} \right] \cdots t_{i,1}^{-1} \right) \\ & \quad \cdot \left(t_{i,1} \cdots t_{i,q_{i+1}} g(a_{i+1}) f(a_{i+1})^{-1} t_{i,q_{i+1}}^{-1} \cdots t_{i,1}^{-1} \right). \end{aligned}$$

As the weight of the representation on the left side of the equality is greater than the weight of the representation on the right side, we have a contradiction. Hence $t_{i,q_{i+1}+1} \neq f(a_{i+1})$.

PROPOSITION 2. *Let $w \in K$ and have yoffi representation $(*)$. Then for each $i \in \{1, 2, \dots, s\}$, $g(a_i)^{-1}t_{i,q_i}^{-1} \cdots t_{i,1}^{-1}t_{i+1,1} \cdots t_{i+1,q_{i+1}}f(a_{i+1})$ in reduced form equals $g_1f_1g_2f_2 \cdots f_{k_i}$ where each $f_j \in F \setminus \{e\}$, $g_j \in G \setminus \{e\}$, and $k_i \geq 1$.*

Proof. It suffices to show that in putting the relevant word in reduced form, the elements $g(a_i)^{-1}$ and $f(a_{i+1})$ do not get cancelled. Indeed, we prove only that $f(a_{i+1})$ does not get cancelled, as the other follows by an analogous argument. The element $f(a_{i+1})$ could get cancelled out only in one of the following ways; but all are impossible, as indicated by the bracketed remarks.

- (1) $t_{i+1,q_{i+1}}^{-1} = f(a_{i+1})$ (Remark 1(i))
- (2) $f(a_{i+1}) = t_{i,q_{i+1}+1}$ and
 $t_{i,j} = t_{(i+1),j}$, $j = 1, \dots, q_{i+1}$. (Lemma 4)
- (3) $t_{i,q_{i+1}}^{-1}t_{(i+1),q_{i+1}}f(a_{i+1}) = e$ and
 $t_{i,j} = t_{(i+1),j}$, $j = 1, \dots, q_{i+1} - 1$ (yoffi implies neutral link).

Thus $f(a_{i+1})$ does not get cancelled out. So we see that the proposition is true.

REMARK 2. Proposition 2 implies that if $w = x_1 \cdots x_s$ is a yoffi representation and has reduced length $\leq n$, then $s \leq n$.

LEMMA 5. *Let $w \in K$ have yoffi representation $(*)$. If the reduced length of $w' = t_{1,1} \cdots t_{1,q_1}v_1 \cdots v_{s-1}t_{(s-1),q_{s-1}}^{-1} \cdots t_{(s-1),1}^{-1}$ is m , then the reduced length of w is greater than or equal to $m - 3$.*

Proof. By Proposition 2, in order to obtain the reduced form of w from the reduced form of w' it suffices to ascertain what is the reduced form of

$$g(a_{s-1})^{-1}t_{s-1,q_{s-1}}^{-1} \cdots t_{s-1,1}^{-1}t_{s,1} \cdots t_{s,q_s}f(a_s)g(a_s)^{-1}t_{s,q_s}^{-1} \cdots t_{s,1}^{-1}.$$

From what we have said earlier it is obvious that the reduced length of $g(a_{s-1})^{-1}t_{s-1,q_{s-1}}^{-1} \cdots t_{s,q_s}f(a_s)$ is greater than or equal to $|q_{s-1} - q_s| - 2$. The reduced length of $g(a_s)t_{s,q_s}^{-1} \cdots t_{s,1}^{-1}$ is greater than or equal to q_s .

Noting that the length of $t_{1,1} \cdots v_1 \cdots f(a_{s-1})$ is greater than or equal to $m - (q_{s-1} + 1)$, we see that the reduced length of w is greater than or equal to $m - q_{s-1} - 1 + |q_{s-1} - q_s| - 2 + q_s \geq m - 3$.

REMARK 3. Of course the bound $m - 3$ in Lemma 5 is a crude one, but it serves the purpose.

PROPOSITION 3. Let $w \in K$ have yoffi representation (*). If the reduced length of w is n then $2q_i + 1 \leq n + 3s$, for each $i = 1, \dots, s$.

Proof. Consider $t_{i,1} \cdots v_i \cdots t_{i,1}^{-1}$. Successively multiplying on the right or left with words of the same kind, we obtain the word w . In so doing, we can apply Lemma 5 above $s - 1$ times, and the result follows.

REMARK 4. Combining Remark 2 and Proposition 3 we obtain the desired result $2q_i + 1 \leq 4n$.

5. Proof of the Theorem. To prove the main result, which is the theorem stated in §2, we must now study the k_ω structure.

It is known [12] that the free product of two k_ω -groups is a k_ω -group. More particularly, if F and G have k_ω decompositions $F = \bigcup_n F_n$ and $G = \bigcup_n G_n$ then the free product has k_ω decomposition

$$F * G = \bigcup_n (F_n \cup G_n)^n.$$

For convenience, we assume $f(A) \subseteq F_1, g(A) \subseteq G_1, F_n = F_n^{-1}, G_n = G_n^{-1}, F_n F_m \subseteq F_{n+m}$ and $G_n G_m \subseteq G_{n+m}$.

Recall that Φ is the canonical continuous homomorphism $F * G \rightarrow F * {}_A G$.

NOTATION. Let

$$X_n = \{ uvu^{-1} : u \in (F_n \cup G_n)^n, v = f(a)g(a)^{-1}, a \in A \}$$

$$X = \bigcup_{n=1}^\infty X_n \quad \text{and} \quad Y_n = (X_n)^n.$$

Here $(X_n)^n$ denotes the set of all words which are the product of at most n elements from X_n .

Clearly each X_n and Y_n is compact, and $K = \bigcup_{n=1}^\infty Y_n$.

REMARK 5. To prove that K is closed, it suffices to show that for each n ,

$$(1) \quad K \cap (F_n \cup G_n)^n \subseteq Y_{2n^3}.$$

This suffices, since then we would have

$$K \cap (F_n \cup G_n)^n = K \cap (F_n \cup G_n)^n \cap Y_{2n^3} = Y_{2n^3} \cap (F_n \cup G_n)^n$$

which is compact, and hence we would have K closed in $F * G$ by the k_ω -condition.

Proposition 1 implies that each $w \in K \cap (F_n \cup G_n)^n$ has a yoffi representation (*). Remarks 2 and 4 then imply that $s \leq n$ and $2q_i + 1 \leq 4n$, for each i . To complete the proof of (1) above (and hence of the theorem) it suffices to show that each $t_{i,j} \in F_{2n^3} \cup G_{2n^3}$. This is a consequence of Proposition 4, proved below.

PROPOSITION 4. *Let $w \in K$ have yoffi representation (*). If $w \in (F_n \cup G_n)^n$, then each $t_{i,j} \in F_{n^3+n} \cup G_{n^3+n}$.*

Proof. By Proposition 2, we know that w in its reduced form is the product of the reduced form of the following blocks:

$$t_{1,1} \cdots t_{1,q_1} f(a_1); \quad g(a_1)^{-1} \cdots f(a_2); \quad g(a_2)^{-1} \cdots f(a_3); \dots;$$

$$g(a_{s-1})^{-1} \cdots f(a_s); \quad g(a_s)^{-1} t_{s,q_s}^{-1} \cdots t_{s,1}^{-1}.$$

The reduced form of w in $F * G$ is unique, so each block in the reduced yoffi representation of w matches a part of the reduced form obtained from the word $w \in (F_n \cup G_n)^n$. Thus any reduced block of w can be expanded to part of a word in $(F_n \cup G_n)^n$ and is therefore itself in $(F_n \cup G_n)^n$.

Consider the block $B_0 = t_{1,1} \cdots t_{1,q_1} f(a_1)$. Recalling that $f(A) \subseteq F_1$, $g(A) \subseteq G_1$, $F_n F_m \subseteq F_{n+m}$ and $G_n G_m \subseteq G_{n+m}$, and noting that each $t_{1,j}$ ($1 \leq j \leq q_1 - 1$) can be expanded to a part of a word in $(F_n \cup G_n)^n$, at worst a product of n elements all from F_n (or G_n) it follows that $t_{1,j} \in F_{n^2} \cup G_{n^2}$. If $t_{1,q_1} \in F$, then, at worst $t_{1,q_1} f(a_1) \in F_{n^2}$, so $t_{1,q_1} \in F_{n^2+1}$. Therefore $t_{1,j} \in F_{n^2+1} \cup G_{n^2+1}$, for $j = 1, \dots, q_1$.

Consider the block $B_1 = g(a_1)^{-1} \cdots f(a_2)$. Each element in B_1 is also in the reduced form (of B_1) or is amalgamated by other elements of B_1 . Such an amalgamation can involve at most three elements. At most one of these three is a $t_{2,j}$. Of the others at most one is a $t_{1,k}$, and the other is in $f(A) \cup g(A)$. As the elements of the reduced form of w lie in $F_{n^2} \cup G_{n^2}$, we see that each element of B_1 must lie in $F_{2n^2+2} \cup G_{2n^2+2}$.

We now proceed to consider in a similar fashion B_2, B_3, \dots, B_{s-1} . We thus obtain that each $t_{i,j}$ that appears in w is an element of $F_{s(n^2+1)} \cup G_{s(n^2+1)}$. The result now follows by observing that $s \leq n$.

REMARK 6. To prove the theorem, it suffices to show that the kernel K of Φ is a closed subgroup of $F * G$. Observing that $\Phi(F) \cap \Phi((F_n \cup G_n)^n) \subseteq \Phi([F_{n^2} \cup (G_{n^2} \cap A)]^n)$ it then follows that F is a closed topological

subgroup of $F *_{\mathcal{A}} G$. Similarly G is seen to be a closed topological subgroup of $F *_{\mathcal{A}} G$ also.

We have thus completed the proof of the theorem.

REMARK 7. We conclude by noting that if G is any connected locally compact group and K is a maximal compact subgroup, then the amalgamated free product $G *_{\mathcal{K}} G$ is seen, from our Theorem, to be a k_{ω} -group (and hence Hausdorff). This example could not be deduced from Ordman [13], Khan and Morris [5] or Katz and Morris [4].

REMARK 8. Our Theorem includes the case when K is a finite group. It should be interesting to investigate this case further.

REFERENCES

- [1] T. H. Fay and B. V. S. Thomas, *Remarks on the free product of two Hausdorff groups*, Arch. Math. (Basel), **33** (1979), 57–65.
- [2] M. I. Graev, *On free products of topological groups*, (Russian), Izv. Akad. Nauk. SSSR, Ser. Mat., **14** (1950), 343–354.
- [3] E. Katz, *Free products in the category of k_{ω} -groups*, Pacific J. Math., **59** (1975), 493–495.
- [4] E. Katz and S. A. Morris, *Free products of k_{ω} -topological groups with normal amalgamation*, Topology Appl., **15** (1983), 189–196.
- [5] M. S. Khan and S. A. Morris, *Free products of topological groups with central amalgamation I*, Trans. Amer. Math. Soc., **273** (1982), 417–432.
- [6] W. F. La Martin, *On the foundations of k -group theory*, Dissertationes mathematicae #146 (1977).
- [7] W. Magnus, A. Karrass and D. Solitar, *Combinatorial Group Theory*, (Dover Publ. Inc., New York, 1976).
- [8] S. A. Morris, E. T. Ordman and H. B. Thompson, *The Topology of Free Products of Topological Groups*, Proc. Second Internat. Conf. Theory of Groups, Canberra (1973), Springer Lecture Notes 372, 504–515.
- [9] B. H. Neumann, *an essay on free products of groups with amalgamation*, Philos. Trans. Royal Soc. London (A), **246** (1954), 503–554.
- [10] P. Nickolas, *A Kurosh subgroup theorem for topological groups*, Proc. London Math. Soc., **42** (81), 461–477.
- [11] E. C. Nummela, *On epimorphisms of topological groups*, General Topology Appl., **9** (1978), 155–167.
- [12] E. T. Ordman, *Free products of topological groups which are k_{ω} -spaces*, Trans. Amer. Math. Soc., **191** (1974), 61–73.
- [13] ———, *Free products of topological groups with equal uniformities I*, Colloq. Math., **31** (1974), 37–43.
- [14] B. V. S. Thomas, *Categories of topological groups*, Memphis State University Report 76-16.
- [15] ———, *On the Coproduct of the Topological Groups Q and Z_2* , Proc. Int. Conf. on categorical topology, Berlin, 1978. Springer Lecture Notes, **719** (1979), 371–375.

Received July 21, 1980 and in revised form July 4, 1984. Some of the research for this paper was carried out while the authors were visiting Tel-Aviv University. The first author thanks La Trobe University for hospitality extended in June to August 1981, and the second author thanks Cleveland State University for hospitality extended in August 1983. The authors gratefully acknowledge the most helpful comments of the referees.

CLEVELAND STATE UNIVERSITY
CLEVELAND, OH 44115

AND

LA TROBE UNIVERSITY
BUNDOORA, VICTORIA
3083, AUSTRALIA