

ON VARIETIES OF TOPOLOGICAL GROUPS
GENERATED BY SOLVABLE GROUPS

BY

SIDNEY A. MORRIS (SYDNEY)

1. In [2] it was shown that any connected Lie group in a variety of topological groups generated by solvable connected Lie groups is solvable. This result is extended here to locally compact groups. (Note that a variety of groups [9] generated by solvable groups may contain non-solvable groups.)

2. A non-empty class of topological groups (not necessarily Hausdorff) is said to be a *variety of topological groups* if it is closed under the operations of taking subgroups, quotients, arbitrary cartesian products and isomorphic images. The smallest variety containing a class \mathcal{A} of topological groups is said to be the *variety generated by \mathcal{A}* , and it is denoted by $V(\mathcal{A})$ (see [1] and [5]-[8]).

If \mathcal{A} is a class of topological groups, then $S\mathcal{A}$ denotes the class of topological groups (topologically) isomorphic to subgroups of members of \mathcal{A} . Similarly we define the operators \bar{S} , \bar{Q} , \mathbf{C} and \mathbf{D} , where they denote closed subgroup, separated quotient, arbitrary cartesian product and finite product, respectively.

We will use the following basic result [1] on generating varieties:

THEOREM. *If \mathcal{A} is a class of topological groups and G is a Hausdorff group in $V(\mathcal{A})$, then $G \in S\mathbf{C}\bar{Q}\bar{S}\mathbf{D}(\mathcal{A})$.*

3. Our first theorem extends the classical result that any connected compact solvable group is abelian.

THEOREM 1. *If \mathcal{A} is a class of solvable Hausdorff groups and G is a connected compact group in $V(\mathcal{A})$, then G is abelian.*

Proof. By the above-mentioned theorem, $G \in S\mathbf{C}\bar{Q}\bar{S}\mathbf{D}(\mathcal{A})$. Since finite products, closed subgroups and separated quotients of solvable topological groups are solvable, this implies that G is a subgroup of a product $\prod_{a \in I} H_a$, where each H_a is solvable. Let $P_a(G)$ be the projection of G into H_a for each $a \in I$. Clearly, each $P_a(G)$ is a connected compact

solvable group and therefore (see 29.44 of [3]) it is abelian. Thus G is a subgroup of a product of abelian groups and hence it is abelian.

THEOREM 2. *If \mathcal{A} is a class of connected locally compact solvable groups and G is a connected locally compact group in $\mathbf{V}(\mathcal{A})$, then G is solvable.*

Proof. Noting Section 4.6 of [4] we see that G has a compact normal subgroup N such that the quotient group G/N is a Lie group. Further, by Section 4.13 of [4], N is contained in a compact connected subgroup M of G . Since $M \in \mathbf{V}(\mathcal{A})$, Theorem 1 shows that M is abelian. Thus N is abelian. We now prove that G/N is solvable and hence G is solvable.

As a consequence of Section 4.6 of [4], we infer that any connected locally compact group H is a subgroup of a product of Lie groups, each of these Lie groups being a quotient of H . This implies that $\mathbf{V}(\mathcal{A}) = \mathbf{V}(\mathcal{B})$, where \mathcal{B} is a class of connected solvable Lie groups. Since G/N is a connected Lie group in $\mathbf{V}(\mathcal{B})$, Corollary 3.4 of [2] implies that G/N is solvable.

REFERENCES

- [1] M. S. Brooks, A. Sidney Morris and Stephen A. Saxon, *Generating varieties of topological groups*, Proceedings of the Edinburgh Mathematical Society (to appear).
- [2] Su-shing Chen and Sidney A. Morris, *Varieties of topological groups generated by Lie groups*, *ibidem* (to appear).
- [3] E. Hewitt and K. A. Ross, *Abstract harmonic analysis II*, Berlin-Heidelberg 1970.
- [4] D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience tracts in pure and applied mathematics I, New York 1955.
- [5] Sidney A. Morris, *Varieties of topological groups*, Bulletin of the Australian Mathematical Society 1 (1969), p. 145-160.
- [6] — *Varieties of topological groups II*, *ibidem* 2 (1970), p. 1-13.
- [7] — *Varieties of topological groups III*, *ibidem* 2 (1970), p. 165-178.
- [8] — *Locally compact abelian groups and the variety of topological groups generated by the reals*, Proceedings of the American Mathematical Society (to appear).
- [9] Hanna Neumann, *Varieties of groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete 37, Berlin-Heidelberg 1967.

THE UNIVERSITY OF NEW SOUTH WALES
KENSINGTON, NEW SOUTH WALES
AUSTRALIA

Reçu par la Rédaction le 24. 7. 1971